

# Examples of relevant ICD-10 codes\* for COSENTYX® (secukinumab) patients

The following ICD-10-CM codes for nr-axSpA are included in the 2021 ICD-10-CM Index to Diseases and Injuries, which goes into effect on October 1, 2020.

Possible nr-axSpA ICD-10-CM Codes	Descriptor
M46.80	Non-radiographic axial spondyloarthritis
M46.82	Non-radiographic axial spondyloarthritis of cervical region
M46.83	Non-radiographic axial spondyloarthritis of cervicothoracic region
M46.84	Non-radiographic axial spondyloarthritis of lumbar region
M46.84	Non-radiographic axial spondyloarthritis of thoracic region
M46.85	Non-radiographic axial spondyloarthritis of thoracolumbar region
M46.87	Non-radiographic axial spondyloarthritis of lumbosacral region
M46.88	Non-radiographic axial spondyloarthritis of sacral and sacrococcygeal region
M46.89	Non-radiographic axial spondyloarthritis of multiple sites

Possible AS ICD-10-CM Codes	Descriptor
M45.0	Ankylosing spondylitis of multiple sites in spine
M45.1	Ankylosing spondylitis of occipito-atlanto-axial region
M45.2	Ankylosing spondylitis of cervical region
M45.3	Ankylosing spondylitis of cervicothoracic region
M45.4	Ankylosing spondylitis of thoracic region
M45.5	Ankylosing spondylitis of thoracolumbar region
M45.6	Ankylosing spondylitis of lumbar region
M45.7	Ankylosing spondylitis of lumbosacral region
M45.8	Ankylosing spondylitis of sacral and sacrococcygeal region
M45.9	Ankylosing spondylitis of unspecified sites in spine

Possible PsA ICD-10-CM Codes	Descriptor
L40.50	Arthropathic psoriasis, unspecified
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.59	Other psoriatic arthropathy

Possible PsO ICD-10-CM Code	Descriptor
L40.0	Plaque psoriasis

AS=ankylosing spondylitis; ICD-10=International Classification of Diseases, 10th Revision; ICD-10-CM=International Classification of Diseases, 10th Revision, Clinical Modification; nr-axSpA=non-radiographic axial spondyloarthritis; PsA=psoriatic arthritis; PsO=plaque psoriasis.

\*These diagnosis code examples are provided for general informational purposes only and are not intended to be directive, a guarantee of coverage, or a substitute for an independent clinical decision.

## INDICATIONS

COSENTYX® (secukinumab) is indicated for the treatment of moderate to severe plaque psoriasis in adult patients who are candidates for systemic therapy or phototherapy.

COSENTYX is indicated for the treatment of adult patients with active psoriatic arthritis.

COSENTYX is indicated for the treatment of adult patients with active ankylosing spondylitis.

COSENTYX is indicated for the treatment of adult patients with active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation.

## IMPORTANT SAFETY INFORMATION

### CONTRAINDICATIONS

COSENTYX is contraindicated in patients with a previous serious hypersensitivity reaction to secukinumab or to any of the excipients.

Please see additional Important Safety Information on back.

Please see full Prescribing Information, including Medication Guide, in pocket.

## **IMPORTANT SAFETY INFORMATION (cont)**

### **WARNINGS AND PRECAUTIONS**

#### **Infections**

COSENTYX may increase the risk of infections. In clinical trials, a higher rate of infections was observed in subjects treated with COSENTYX compared to placebo-treated subjects. In placebo-controlled clinical trials in patients with moderate to severe plaque psoriasis, higher rates of common infections such as nasopharyngitis (11.4% versus 8.6%), upper respiratory tract infection (2.5% versus 0.7%), and mucocutaneous infections with candida (1.2% versus 0.3%) were observed with COSENTYX compared with placebo. A similar increase in risk of infection was seen in placebo-controlled trials in patients with psoriatic arthritis, ankylosing spondylitis and non-radiographic axial spondyloarthritis. The incidence of some types of infections appeared to be dose-dependent in clinical studies.

Exercise caution when considering the use of COSENTYX in patients with a chronic infection or a history of recurrent infection.

Instruct patients to seek medical advice if signs or symptoms suggestive of an infection occur. If a patient develops a serious infection, the patient should be closely monitored and COSENTYX should be discontinued until the infection resolves.

#### **Pre-treatment Evaluation for Tuberculosis**

Evaluate patients for tuberculosis (TB) infection prior to initiating treatment with COSENTYX. Do not administer COSENTYX to patients with active TB infection. Initiate treatment of latent TB prior to administering COSENTYX. Consider anti-TB therapy prior to initiation of COSENTYX in patients with a past history of latent or active TB in whom an adequate course of treatment cannot be confirmed. Patients receiving COSENTYX should be monitored closely for signs and symptoms of active TB during and after treatment.

#### **Inflammatory Bowel Disease**

Caution should be used when prescribing COSENTYX to patients with inflammatory bowel disease. Exacerbations, in some cases serious, occurred in patients treated with COSENTYX during clinical trials in plaque psoriasis, psoriatic arthritis, ankylosing spondylitis and non-radiographic axial spondyloarthritis. In addition, new onset inflammatory bowel disease cases occurred in clinical trials with COSENTYX. In an exploratory study in 59 patients with active Crohn's disease, there were trends toward greater disease activity and increased adverse events in the secukinumab group as compared to the placebo group. Patients who are treated with COSENTYX should be monitored for signs and symptoms of inflammatory bowel disease.

#### **Hypersensitivity Reactions**

Anaphylaxis and cases of urticaria occurred in patients treated with COSENTYX in clinical trials. If an anaphylactic or other serious allergic reaction occurs, administration of COSENTYX should be discontinued immediately and appropriate therapy initiated.

The removable cap of the COSENTYX Sensoready® pen and the COSENTYX prefilled syringe contains natural rubber latex which may cause an allergic reaction in latex-sensitive individuals. The safe use of the COSENTYX Sensoready pen or prefilled syringe in latex-sensitive individuals has not been studied.

#### **Vaccinations**

Prior to initiating therapy with COSENTYX, consider completion of all age appropriate immunizations according to current immunization guidelines. Patients treated with COSENTYX should not receive live vaccines.

Non-live vaccinations received during a course of COSENTYX may not elicit an immune response sufficient to prevent disease.

### **MOST COMMON ADVERSE REACTIONS**

Most common adverse reactions (>1%) are nasopharyngitis, diarrhea, and upper respiratory tract infection.

**Please see additional Important Safety Information on front.**

**Please see full Prescribing Information, including Medication Guide, in pocket.**

